

Product Information

Anti-HOXB13

produced in rabbit, affinity isolated antibody

Catalog Number **H1416**

Product Description

Anti- HOXB13 is produced in rabbit using as immunogen a synthetic peptide corresponding to amino acids 270-284 of human HOXB13 (GeneID: 10481) conjugated to KLH. The corresponding sequence differs by 2 amino acids in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-HOXB13 recognizes human HOXB13. The antibody may be used in several immunochemical techniques including immunoblotting (~30 kDa) and immunoprecipitation. Detection of the HOXB13 band by immunoblotting is specifically inhibited with the immunizing peptide.

Hox genes are evolutionarily conserved transcription factors, which act to control important development pathways involved in morphogenesis of the embryo. In vertebrates, there are 39 HOX genes that are organized into four clusters (*HOXA*–*HOXD*) located on different chromosomes. Each cluster contains 9–11 member genes encoding relatively small gene products, all containing a highly conserved 60-amino-acid region (the homeobox) with DNA-binding activity that contributes to their activity as transcription factors.¹ One of the major functions of *Hox* genes seems to be the formation of the body plan during embryonic development.² In addition to roles in normal development, altered homeobox gene function or expression is implicated in the development of cancers, such as leukemias or in neoplasms of the breast, prostate, kidney, colon, skin and brain.^{3,4} HOXB13 is overexpressed in human endometrial, ovarian, and cervical carcinomas and is associated with the invasiveness of ovarian and endometrial cancer cells.^{5–7}

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2–8 °C for up to one month. For extended storage, freeze in working aliquots at –20 °C. Repeated freezing and thawing, or storage in “frost-free” freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 2–4 µg/mL is recommended using extracts of HEK-293T cells.

Immunoprecipitation: a working antibody amount of 2.5–5 µg is recommended using lysates of HeLa cells.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Lemons, D., et al., *Science*, **313**, 1918–1922 (2006).
2. Akam, M., *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **349**, 313–319 (1995).
3. Stuart, E.T., et al., *Adv. Genet.*, **33**, 255–274 (1995).
4. Cillo, C., et al., *Exp. Cell Res.*, **248**, 1–9 (1999).
5. Lopez, R., et al., *Int. J. Gynecol. Cancer*, **16**, 329–335 (2006).
6. Yamashita, T., et al., *Int. J. Oncol.*, **28**, 931–938 (2006).
7. Zhao, Y., et al., *Oncol. Rep.*, **13**, 721–726 (2005).

SG,KAA,PHC 05/09-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.