SigmaAldrich.com

Sigma-Aldrich.

Product Information

Anti-FAM134B (Lumenal) Antibody

Produced in Rabbit Affinity Isolated Antibody

SAB4200848

Product Description

Anti-FAM134B (lumenal) antibody is developed in rabbit using synthetic peptide corresponding to the internal region of human FAM134B (GeneID: 54463), conjugated to KLH as immunogen. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-FAM134B (lumenal) antibody specifically recognizes an epitope on human FAM134B ER luminal domain. The antibody may be used in various immunochemical techniques including immunoblotting (~ 55 kDa) and immunofluorescence. Detection of the FAM134B band by immunoblotting is specifically inhibited by the immunogen.

Five endoplasmic reticulum (ER) resident proteins have been shown to function as receptors for selective ER-phagy: FAM134B, SEC62, RTN3, CCPG1 and ATL3.¹⁻² These integral membrane proteins appear to partition into different sub-domains of the ER. All of them harbor a sequential peptide motif within the cytoplasmic region, enabling binding to LC3/GABARAP proteins associated with phagophore membrane.¹ FAM134B (Synonyms: RETREG1 or JK-1) is the first ER-phagy receptor to be identified.² The protein was discovered to be involved in the pathogenesis of esophageal squamous cell carcinoma (ESCC).³⁻⁴ Loss of function mutations in FAM134B result in severe sensory and autonomic neuropathy (HSANII).⁵ In addition, FAM134B has been classified as an intra-membrane ER-resident protein that is characterized by the presence of a reticulon homology domain (RHD) and that is mainly located at the edges of the ER sheets.^{1,6} In research, FAM134B is implicated in the suppression of viral replication during Ebola, Dengue, Zika and West Nile viral infections.7-8

Non-structural viral proteases, such as NS2S3, specifically cleave FAM134B, thereby subverting ER-phagy.⁷ Specific mutations in FAM134B and expression profiles are strongly correlated with colorectal cancers.⁹ Increased expression of FAM134B is also implicated with susceptibility to vascular dementia and allergic rhinitis.⁹⁻¹⁰

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody Concentration: ~ 1.0 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working dilution of 1:500-1:1000 is recommended using whole extract of HEK-293 cells over-expressing human FAM134B.



Immunofluorescence: a working dilution of 1:500-1:1000 is recommended using human HeLa cells.

Note: In order to obtain best results in different techniques and preparations it is recommended to determine optimal working concentration by titration test.

References

- Bhaskara RM., et al., *Nat Commun.*, **10**: 2370 (2019).
- 2. Khaminets A., et al., *Nature*, **522**: 354-8 (2015).
- 3. Tang JC., et al., *Clin Cancer Res.*, **7**: 1539-45 (2001).
- Tang WK., et al., Int J Mol Med., 19: 915-23 (2007).
- 5. Kurth I., et al., Nat Genet., 41: 1179-81 (2009).
- 6. Grumati P., et al., J Cell Sci., 131: (2018).
- Lennemann NJ. and Coyne CB., Autophagy, 13: 322-32 (2017).
- Chiramel AI., et al., J Infect Dis., 214: S319-25 (2016).
- Islam F., et al., Genes Chromosom Cancer, 57: 240-51 (2018).
- 10. Melchiotti R., et al., *BMC Med Genet.*, **15**: 73 (2014).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at <u>SigmaAldrich.com/techservice</u>.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at <u>SigmaAldrich.com/terms</u>.

Contact Information

For the location of the office nearest you, go to <u>SigmaAldrich.com/offices</u>.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. SAB4200848dat Rev 06/21

