

Product Information

Anti-IRAK2, C-Terminal

produced in rabbit, IgG fraction of antiserum

Catalog Number **I1529**

Synonym: Anti-Interleukin-1 Receptor-Associated Kinase 2

Product Description

Anti-IRAK2, C-Terminal is produced in rabbit using a synthetic peptide corresponding to the C-terminal amino acids 571-590 of human IL-1 receptor-associated kinase 2 (IRAK2)¹ as immunogen.

Anti-IRAK2, C-Terminal recognizes human IRAK2 (65 kDa) by immunoblotting using a K562 cell lysate. No cross-reactivity is observed with IRAK.

A novel member of the IRAK/Pelle family has been identified and designated IRAK2. Members of this family play a central role in IL-1R1 and Toll-Like Receptor (TLR) mediated inflammatory/immune response, leading to the activation of NF- κ B.² Binding of the pro-inflammatory cytokine IL-1 to its receptors (IL-1R) leads to the association of IL-1R with the IL-1 receptor accessory protein (IL-1RAcP).^{3,4} An adapter protein, myeloid differentiation protein (MyD88), is then recruited to this complex and activated.^{5,6} MyD88 then recruits to the complex, two distinct putative serine-threonine kinases: IL-1 receptor-activated kinase (IRAK) and IRAK2.⁷ These kinases then interact with the adapter molecule, TNF receptor-associated factor 6 (TRAF6),^{8,9} which links them to the protein kinase NF κ B-inducing kinase (NIK).¹⁰ NIK activates the I κ B kinase complex (IKK α and IKK β) that directly phosphorylates I κ B.¹¹⁻¹³ The phosphorylation of I κ B initiates ubiquitin-proteasome-mediated degradation which subsequently activates transcription factor NF- κ B.^{14,15}

IRAKs are also known to associate with Toll like receptor (TLR) that plays a critical role in innate immunity against bacteria, and the dominant negative mutants of IRAKs inhibit lipopolysaccharide (LPS) induced NF- κ B activation.^{16,17} IRAK2 is expressed in a variety of human tissues.

Reagents

Supplied at ~0.5 mg/ml in phosphate buffered saline, containing 0.02% sodium azide.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Antibody can be stored at 2-8 °C for three months and at -20 °C for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Product Profile

Immunoblotting: the recommended working concentration is 0.5-1 μ g/ml using a K562 cell lysate.

References

1. Muzio, M., et al., *Science*, **278**, 1612-1615 (1997).
2. Cao, Z., et al., *Science*, **271**, 1128-1131 (1996).
3. Huang, J., et al., *Proc. Natl. Acad. Sci. USA*, **94**, 12829-12832 (1997).
4. Croston, G. E., et al., *J. Biol. Chem.*, **270**, 16514-16517 (1995).
5. Wesche, H., et al., *Immunity*, **7**, 837-847 (1997).
6. Burns, K., et al., *J. Biol. Chem.*, **273**, 12203-12209 (1998).
7. Muzio, M., et al., *Science* **278**, 1612-1615 (1997).
8. Cao, Z., et al., *Science* **383**, 443-446 (1996).
9. Muzio, M., et al., *J. Exp. Med.* **187**, 2097-2101 (1998).
10. Song, H. Y., et al., *Proc. Natl. Acad. Sci. USA*, **94**, 9792-9796 (1997).
11. DiDonato, J. A., et al., *Nature*, **388**, 548-554 (1997).
12. Stancovski, I., and Baltimore, D. *Cell*, **91**, 299-302 (1997).
13. Scheidereit, C. *Nature*, **395**, 225-226 (1998).
14. Baeuerle, P. A., and Baltimore, D. *Cell*, **87**, 13-20 (1996).
15. Baeuerle, P. A. *Curr. Biol.*, **8**, R19-R22 (1998).
16. Zhang, F.X., et al., *J. Biol. Chem.*, **274**, 7611-7614 (1999).
17. Yang, R.B., et al., *J. Immunol.*, **163**, 639-643 (1999).

RC,PHC 09/12-1