

Product Information

Anti-EGF Receptor Antibody, Mouse Monoclonal

Clone 225, purified from hybridoma cell culture

E2156

Product Description

Anti-EGFR (mouse IgG1 isotype) is derived from the hybridoma 225 produced by the fusion of mouse myeloma cells (NS-1-503 cells) and splenocytes from BALB/c mice immunized with partially purified EGF receptors from A-431 cells. The isotype is determined using a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents, Cat. No.ISO2.

Anti-EGF Receptor recognizes human EGF Receptor. The antibody blocks the binding of EGF to the receptor, inhibits the growth of A-431 tumor cells *in vitro*,¹⁻² and activates caspase-8 thus inducing apoptosis in DiFi colon cancer cells.³ The antibody can be used in immunoprecipitation.

The receptor for epidermal growth factor is an integral cell membrane glycoprotein of 170 kDa, which spans the membranes of a wide range of normal and malignant epithelial cells.⁴⁻⁵ The EGF receptor has an intracellular domain that exhibits tyrosine kinase activity. Overexpression of the receptor can produce a neoplastic phenotype in cells, which has been associated with poor prognosis or short survival in patients with several types of cancer. EGFR protein tyrosine kinase is activated when EGF binds the extracellular binding domain. The first detectable response is the autophosphorylation of the C-terminal tyrosines followed by phosphorylation of other endogenous substrates. The phosphorylated sequences of EGFR have a high affinity to SH2 domain containing proteins. These proteins are phosphorylated and activated by the receptor tyrosine kinases.⁶ Activation of EGFR can be achieved in the absence of EGF by receptor dimerization with antibody⁷ or by 10 mM Mn²⁺ ion.

Reagent

Supplied in 0.01 M phosphate buffered saline, pH 7.4, 0.2 µm filtered.

Antibody concentration: ~1.5 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoprecipitation

4-8 µg of antibody can immunoprecipitate EGF Receptor from A431 cell lysate.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Sato, J.D., et al., Mol. Biol. Med., 1, 511-529 (1983).
2. Masui, H., et al., Cancer Res., 44, 1002-1007 (1984).
3. Liu, B., and Fan, Z., Oncogene, 20, 3726-3734 (2001).
4. Ullrich, A., et al., Nature, 309, 418-425 (1984).
5. Carpenter, G., Annu. Rev. Biochem., 56, 881-914 (1987).
6. Margolis, B., et al., EMBO J., 9, 4375-4380 (1990).
7. Yarden, Y., and Schlessinger, J., Biochemistry, 26, 1443-1451 (1987).

This product is for in vitro research use only. It is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
E2156dat Rev 05/22

The Merck logo, consisting of the word "MERCK" in a bold, red, sans-serif font.