

## Product Information

### CHICKEN IgG

Purified Immunoglobulin  
Reagent Grade

Product No. **I 4881**

#### Product Description

Purified Chicken IgG is isolated from pooled normal chicken serum by fractionation and ion-exchange chromatography. The product is lyophilized from 0.01 M phosphate buffer, pH 7.2.

#### Uses

Purified IgG may be used as a reference antigen, standard, blocking agent or coating protein in a variety of immunoassays including ELISA, dot immunobinding, Western immunoblotting, immunodiffusion, and immunoelectrophoresis. Other applications include starting materials for the preparation of immunogens and solid-phase immunoadsorbents.

#### Reconstitution

The product should be reconstituted to a minimum of 1 mg/ml with a saline (150 mM NaCl) solution. Protein concentration should be determined after reconstitution by absorbance at 280 nm ( $E_{280}^{1\%} = 14$ ).

#### Storage

Prior to reconstitution, store at 2-8 °C. After reconstitution, store aliquots at -20 °C. Repeated freezing and thawing is **not** recommended.

#### Performance Characteristics

Appearance: White, lyophilized powder.

Solubility: Slightly hazy, colorless solution at 50 mg/ml in 0.85% NaCl.

Sodium Content (ICP-Atomic Emission): <10%

Water Content (Karl Fischer): <10%

Identity (Immunoelectrophoresis): Single arc of precipitation in the gamma region at 50 and 250 µg of immunoglobulin versus anti-chicken whole serum and anti-chicken IgG.

Purity (SDS-PAGE): Not less than 95%.

10/98

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.