sigma-aldrich.com

3050 Spruce Street, Saint Louis, MO 63103 USA

Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757 email: techservice@sial.com

Product Information

Anti-CDC25c antibody produced in chicken, affinity isolated antibody

Catalog Number **GW21532** Formerly listed as GenWay Catalog Number 15-288-21532, M-phase inducer phosphatase 3 Antibody.

Storage Temperature Store at −20 °C

Synonyms: Cell division cycle 25C protein isoform b, EC 3.1.3.48; Dual specificity phosphatase Cdc25C

Product Description

Functions as a dosage-dependent inducer in mitotic control. It is a tyrosine protein phosphatase required for progression of the cell cycle. It directly dephosphorylates CDC2 and activate its kinase activity.

NCBI Accession number: NP_073720.1 Swiss Prot Accession number: P30307

Gene Information: Human .. CDC25C (995)

Immunogen: Recombinant protein Cell division cycle 25C

protein isoform b

Immunogen Sequence: GI # 12408658, sequence 351 - 400

The product is a clear, colorless solution in phosphate buffered saline, pH 7.2, containing 0.02% sodium azide.

Species Reactivity: Human

Tested Applications: WB

Recommended Dilutions: Recommended starting dilution for Western blot analysis is 1:500, for tissue or cell staining 1:200.

<u>Note</u>: Optimal concentrations and conditions for each application should be determined by the user.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Due to the sodium azide content a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2–8 °C for up to one week. For extended storage, store in –20 °C freezer in working aliquots. Repeated freezing and thawing, or storage in "frostfree" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

LPG.MAM12/09-1