



**ANTI-PHOSPHO-SEROTONIN  
N-ACETYLTRANSFERASE (C-TERMINAL) (pSer<sup>206</sup>)**  
Developed in Rabbit, Affinity Isolated Antibody

Product Number **S 0939**

Anti-phospho-Serotonin N-Acetyltransferase (C-terminal) (pSer<sup>206</sup>) is developed in rabbit using as immunogen a synthetic phosphopeptide derived from the C-terminus of the rat arylalkylamine N-acetyltransferase (AA-NAT) containing serine 206. The serum is affinity purified using epitope-specific affinity chromatography. Anti-phospho-AA-NAT (pSer<sup>206</sup>) recognizes rat AA-NAT (pSer<sup>206</sup>). It has been used in immunoblotting applications.

Pineal gland serotonin AA-NAT, a 23 kDa protein, is the rate-limiting enzyme in pineal and retinal melatonin synthesis. The nocturnal oscillation of enzyme activity ranges from 10- to 100- fold depending on vertebrate species. The enzyme converts serotonin to N-acetylserotonin.<sup>1,2</sup>

Studies have shown that AA-NAT is regulated by cAMP at both mRNA and protein levels. Phosphorylation of AA-NAT promotes formation of a complex AA-NAT/14-3-3 which protects AA-NAT from dephosphorylation and increases melatonin production.<sup>3,4</sup> AA-NAT has two protein kinase A binding sites: threonine 29 at N-terminal and serine 206 at C-terminal. Phosphorylation of AA-NAT at these residues might play a role in protein inactivation by proteasome-dependent degradation.

**Reagent**

Anti-phospho-Serotonin N-Acetyltransferase (N-terminal) (pSer<sup>206</sup>) is supplied as 50 µg of purified rabbit IgG in phosphate buffered saline (PBS) containing 1 mg/ml BSA and 0.05% sodium azide

## Product Information

**Precautions and Disclaimer**

Due to the sodium azide content a material safety sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

**Storage/Stability**

Store at -20 °C. For extended storage, upon initial thawing, freeze in working aliquots. Avoid repeated freezing and thawing to prevent denaturing the antibody. Working dilution samples should be discarded if not used within 12 hours. The antibody is stable for at least 6 months when stored appropriately. Do not store in a frost-free freezer.

**Product Profile**

A recommended working concentration of 0.5 µg/ml is determined by immunoblotting using cell lysates from transfected 293T cells. Data demonstrate that only phosphopeptide corresponding to the region containing serine 206 blocks the antibody signal, confirming the specificity of Anti-phospho-Serotonin N-Acetyltransferase (N-terminal) (pSer<sup>206</sup>) for this phosphos phosphorylated residue

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working concentration by titration test.

**References**

1. Zhan-Poe, X. and Craft, C.M., J. Pineal. Res., **27**, 49-58 (1999).
2. Herichova, I. et al., Neurosci. Lett., **298**, 123-126 (2001).
3. Li, X., et al., Curr. Opin. Neurobiol., **8**, 648-651 (1998).
4. Ganguly, S., et al., Proc. Natl. Acad. Sci. USA, **98**, 8083-8088 (2001).

AH 7/01

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.