

## Ecteola Cellulose

Product Number **E9251**  
Store at Room Temperature

### Product Description

Ecteola cellulose is prepared by activating cellulose with epichlorohydrin, followed by reacting the epichlorohydrin-activated cellulose with triethanolamine.

Ecteola cellulose has been used for removal of heparin from plasma.<sup>1,2</sup> Heparinase is, however, more effective than ecteola cellulose in removing heparin from plasma.<sup>1</sup>

Several naturally occurring polysaccharides were purified, subsequently sulfated by chlorosulfonic acid-pyridine complex, isolated as the sodium salt, and further purified by ecteola cellulose chromatography. The anticoagulant properties of the sulfated polysaccharides were then studied.<sup>3</sup>

Ecteola cellulose has been used in the purification of a sulfated trisaccharide,<sup>4</sup> and in the purification of bovine lung and heart glycosaminoglycans (GAGs).<sup>5</sup> It has been used in an assay procedure for the enzyme 3'-phosphoadenosine 5'-phosphosulfate:phenol sulfotransferase.<sup>6</sup> It has also been used in the separation of purine analogs by thin layer chromatography (TLC).<sup>7</sup>

### Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

### References

1. Van den Besselaar, A. M. and Meeuwisse-Braun, J., Enzymatic elimination of heparin from plasma for activated partial thromboplastin time and prothrombin time testing. *Blood Coagul. Fibrinolysis*, **4(4)**, 635–638 (1993).
2. McManus, M. L., et al., Coagulation factor deficiencies during initiation of extracorporeal membrane oxygenation. *J. Pediatr.* **126(6)**, 900–904 (1995).
3. Doctor, V. M., et al., Anticoagulant properties of semisynthetic polysaccharide sulfates. *Thromb. Res.* **64(4)**, 413–425 (1991).
4. Skelton, T. P., et al., Characterization of a sulfotransferase responsible for the 4-O-sulfation of terminal beta-N-acetyl-D-galactosamine on asparagine-linked oligosaccharides of glycoprotein hormones. *J. Biol. Chem.* **266(26)**, 17142–17150 (1991).
5. Volpi, N., Disaccharide analysis and molecular mass determination to microgram level of single sulfated glycosaminoglycan species in mixtures following agarose-gel electrophoresis. *Anal Biochem.* **273(2)**, 229–239 (1999).
6. Borchardt, R. T. et al., An Ecteola-cellulose chromatography assay for 3'-phosphoadenosine 5'-phosphosulfate: phenol sulfotransferase. *Anal Biochem.* **130(2)**, 334–338 (1983).
7. Harber, M. J. and J. L. Maddocks, Separation of purine analogues from nucleic acid derivatives on a thin layer of ecteola-cellulose. *J. Chromatogr.* **101**, 231–234 (1974).

IRB 11/07

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.