

Technical Data Sheet

Buffered NaCl-Peptone Solution + 0.1 % Tween® 80

Ordering number: 1.46612.0010

Buffered NaCl-Peptone Solution with 0.1 % Tween® 80 is designed for the microbial examination of non-sterile pharmaceutical products according to the European Pharmacopoeia and the United States Pharmacopoeia.

The solution is used for the microbial examination of non-sterile pharmaceutical products, e.g. as solution for sample preparation or as dilution or washing solution.

The formulation is prepared according to the recommendations of the European Pharmacopoeia, United States Pharmacopoeia and Japanese Pharmacopoeia (EP, 2.6.13.; JP, 5.02 and USP, 62).

The Buffered NaCl-Peptone Solution is available with different Tween® 80 concentration, in different filling volumes and with various locking mechanisms:

- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146164): 17 ml-tube, filling volume 9 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146371): 275 ml-bottle with **wide mouth**, filling volume 90 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146612): 125 ml-bottle with **flip cap**, filling volume 100 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146421): 250 ml-bottle with **screw cap**, filling volume 200 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146634): 500 ml-bottle with **screw cap**, filling volume 450 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146381): 1000 ml-bottle with **screw cap**, filling volume 1000 ml
- Buffered NaCl-Peptone Solution + 0.1 % Tween® 80 (article number 146633): 1000 ml-bottle with **flip cap**, filling volume 1000 ml
- Buffered NaCl-Peptone Solution + 1% Tween® 80 (article number 146698): 500 ml-bottle with **screw cap**, filling volume 500 ml
- Buffered NaCl-Peptone Solution + 1% Tween® 80 (article number 146651): 1000 ml-bottle with **screw cap**, filling volume 1000 ml

Mode of Action

The low content of peptone as well as the content of phosphate buffer and sodium chloride in the solution maintains the viability of present microorganisms without supporting growth. The addition of polysorbate 80 (Tween® 80, which is a registered trademark of ICI Americas Inc.) may assist the suspension of poorly wettable substances. It may further serve as neutralizing agent for disinfectants such as quaternary ammonium compounds, iodines and parabenes.

Typical Composition

Additive	Buffered NaCl + 0.1 % Tween®80	Buffered NaCl + 1 % Tween®80
Peptone	1 g/l	1 g/l
NaCl	4.3 g/l	4.3 g/l
KH ₂ PO ₄	3.6 g/l	3.6 g/l
Na ₂ HPO ₄ · 2 H ₂ O	7.2 g/l	7.2 g/l
Polysorbate (Tween®) 80	1 ml/l	10 ml/l

The appearance of the medium is clear and colorless. The pH value is in the range of 6.8-7.2. The medium can be adjusted and/or supplemented according to the performance criteria required.

Application and Interpretation

For quality control a stability test is performed in order to test the ability of the solution to maintain viability of microorganisms. Therefore the buffer solution to be tested is inoculated with 100-1,000 CFU/ml. Directly after inoculation as well as after an incubation time of 1 hour at 20 to 25 °C for each test strain 100 µl are subcultured on Columbia Blood Agar (article number 146559). The viable count should not change from the beginning to the end of incubation. A typical stability test is shown under "Quality Control".

Storage and Shelf Life

The product can be used for tests until the expiry date if protected from light and properly sealed at +2 °C to +25 °C.

The testing procedures as described on the CoA can be started up to the expiry date printed on the label.

Disposal

Please mind the respective regulations for the disposal of used culture medium (e.g. autoclave for 20 min at 121 °C, disinfect, incinerate etc.).



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

Merck Millipore and the M logo are registered trademarks of Merck KGaA, Darmstadt, Germany. Lit. No. TN7716EN00

Quality Control

Control Strains	ATCC #	Inoculum CFU	Incubation	Expected Results
<i>Staphylococcus aureus</i>	6538	2,000-20,000	1 h ± 5 min at 20-25 °C	No change in the number of CFU
<i>Pseudomonas aeruginosa</i>	9027	2,000-20,000	1 h ± 5 min at 20-25 °C	No change in the number of CFU
<i>Candida albicans</i>	10231	2,000-20,000	1 h ± 5 min at 20-25 °C	No change in the number of CFU
<i>Escherichia coli</i>	8739	2,000-20,000	1 h ± 5 min at 20-25 °C	No change in the number of CFU
<i>Bacillus subtilis</i>	6633	2,000-20,000	1 h ± 5 min at 20-25 °C	No change in the number of CFU

Please refer to the actual batch related Certificate of Analysis.

Literature

European Pharmacopoeia 8.0 (2014): 2.6.13. Microbial examination of non-sterile products (test for specified microorganisms).

Japanese Pharmacopoeia 16th edition (2011): 5.02 Microbial Limit Test for Crude Drugs.

United State Pharmacopoeia 39 NF 34 (2015): <62> Microbial examination of non-sterile products (test for specified microorganisms).

Ordering Information

Product	Cat. No.	Pack size
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46164.0100	100 x 9 ml tube
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46371.0006	6 x 90 ml bottle
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46612.0010	10 x 100 ml bottle
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46421.0006	6 x 200 ml bottle
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46634.0006	6 x 450 ml bottle
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46381.0006	6 x 1000 ml bottle
Buffered NaCl-Peptone Solution + 0.1 % Tween® 80	1.46633.0006	6 x 1000 ml bottle
Buffered NaCl-Peptone Solution + 1% Polsorb. 80	1.46681.0006	6 x 350 ml bottles
Buffered NaCl-Peptone Solution + 1% Polsorb. 80	1.46698.0006	6 x 500 ml bottles
Buffered NaCl-Peptone Solution + 1% Polsorb. 80	1.46651.0006	6 x 1000 ml bottles
Buffered NaCl-Peptone Solution + 5% Polsorb. 80	1.46682.0010	10 x 20 ml bottles
Columbia Blood Agar Pharm.	1.46559.0020	20 x 90mm plates
Columbia Blood Agar Pharm.	1.46559.0100	100 x 90mm plates

Merck KGaA, 64271 Darmstadt, Germany
Fax: +49 (0) 61 51 / 72-60 80
mibio@merckgroup.com
www.merckmillipore.com/biomonitoring

Find contact information for your country at:
www.merckmillipore.com/offices
For Technical Service, please visit:
www.merckmillipore.com/techservice



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

Merck Millipore and the M logo are registered trademarks of Merck KGaA, Darmstadt, Germany. Lit. No. TN7716EN00