



SIGMA-ALDRICH

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

Collagenase from *Clostridium histolyticum*

Product Number **C1889**
Storage Temperature -0 °C

Product Description

Enzyme Commission (EC) Number: 3.4.24.3
CAS Number: 9001-12-1

This collagenase is obtained from the culture filtrate of *Clostridium histolyticum*. The culture filtrate is thought to contain at least 7 different proteases ranging in molecular weight from 68-130 kDa.¹ This lyophilized powder is prepared from a sterile filtered solution of Product No. C 5138.

Collagenase is typically used to digest the connective components in tissue samples to liberate individual cells. The concentration for cartilage dispersal is 1-2 mg/ml, but literature searches should be performed for species specific and/or tissue specific concentrations.^{2,3}

Many references exist for using collagenase to digest various tissues. The choice of one technique over another is often arbitrary and based more on past experience than on an understanding of why the method works and what modifications could lead to better results. Concentrations typically vary from 0.1 to 5 mg/ml, and digestion time should be experimentally monitored using a very gentle agitation system to check for tissue dissociation. Collagenase treatment can cause some cells to die. Satisfactory efficiency of cell dissociation without causing too much cell death typically is achieved from 15 minutes to several hours, but can fall outside of this range if the concentration is unusual. The preferred buffer to use is Krebs Ringer Buffer with calcium and BSA. Zn^{2+} is required for activity, but it is tightly bound to the collagenase during purification. Additional Zn^{2+} should not be necessary as long as no chelator is added to the solution during digestion.

When this enzyme is tested for suitability for the release of hepatocytes, the collagenase is used at approximately 1 mg/ml in a total volume of 100 ml for each rat liver.

If excessive cell death is observed with concentrations used with previous lots, the new lot used might have a higher specific activity. Lowering the enzyme concentration and/or adding BSA or serum (0.5% and 5-10%, respectively) is recommended. These components are added to stabilize the cells to further digestion by the enzyme.

Radiolabeled gelatin has been used to measure the activity and mechanism of collagenase digestion.⁴

Mandl units have the same description as Sigma collagen digestion units. The conversion factor for Mandl units/Wuensch units to Sigma units is approximately 1000-2000 to 1.

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

For measurement of enzymatic activity, an enzyme stock solution is prepared by dissolving 0.05 - 0.1 mg/ml collagenase in 50 mM TES buffer, pH 7.4 (37 °C), containing 0.36 mM calcium chloride. Final concentrations in the reaction mixture are 50 mM TES, 0.36 mM calcium chloride, 25 mg collagen (Product No. C9879), and 0.005-0.01 mg collagenase.

For tissue culture applications, collagenase can be solubilized in calcium-free solutions such as Hank's Balanced Salts (Product No. H2387) or Earle's Balanced Salt Solution (Product No. E6267).

To sterile filter solutions of collagenase, first centrifuge the solution or filter through a 0.8 µm filter to remove insolubles. This will remove particulates and reduce the probability of clogging the 0.2 µm filter during sterile filtration.

Storage/Stability

Solutions at neutral pH and with adequate calcium ion concentration (0.3-0.5 mM) will retain activity for at least 5 hours at 37 °C.

Solutions at -20 °C are stable for several months.³

References

1. Angleton, E.L., et. al., Preparation and reconstitution with divalent metal ions of class I and class II *Clostridium histolyticum* apocollagenases. *Biochemistry*, **27**, 7406-7412 (1988).
2. Bassleer, C., et al., Human Chondrocytes in Tridimensional Culture, *In Vitro Cell. Dev. Biol.*, **22**, 113-119 (1986).
3. Klagsbrun, M., Large-Scale Preparation of Chondrocytes, *Methods in Enzymology*, **58**, 560-564 (1979).
4. Mookhtiar, K. A., et al., Properties of Radiolabeled Type I, II, and III Collagens Related to their Use as Substrates in Collagenase assays, *Anal. Biochem.*, **158**, 322-333 (1986).

MWM/RXR 5/06

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.