

C3097 Campylobacter Supplement II (Butzler)

An antibiotic supplement recommended for the selective isolation of the Campylobacter species.

Composition

(per vial, sufficient for 500 ml medium)

Bacitracin 1'2500 IU

Colistin Sulphate 5'000 units

Cycloheximide 25 mg

Cephazolin Sodium 7.5 mg

Novobiocin 2.5 mg

Store at 4°C.

Directions

Rehydrate the contents of one vial aseptically with 5 ml of 50% ethanol and mix well to dissolve. Avoid frothing of the solution. Aseptically, add the contents of the vial to 500 ml of sterile, molten Columbia Agar Base (Fluka 27688). Mix gently and pour into sterile petri plates.

C3097 Campylobacter Supplement II (Butzler)

An antibiotic supplement recommended for the selective isolation of the Campylobacter species.

Composition

(per vial, sufficient for 500 ml medium)

Bacitracin 1'2500 IU

Colistin Sulphate 5'000 units

Cycloheximide 25 mg

Cephazolin Sodium 7.5 mg

Novobiocin 2.5 mg

Store at 4°C.

Directions

Rehydrate the contents of one vial aseptically with 5 ml of 50% ethanol and mix well to dissolve. Avoid frothing of the solution. Aseptically, add the contents of the vial to 500 ml of sterile, molten Columbia Agar Base (Fluka 27688). Mix gently and pour into sterile petri plates.