



Product Information

Manganese(II) sulfate monohydrate Cell Culture Tested

Product Number **M 1144**
Store at Room Temperature

Product Description

Molecular Formula: $\text{MnSO}_4 \cdot \text{H}_2\text{O}$
Molecular Weight: 169.0
CAS Number: 10034-96-5

This product is cell culture tested (0.15 ng/ml) and is appropriate for use in cell culture experiments.

Manganese sulfate occurs in nature as several mineral forms, jokokuite, pentahydrate, szmikite, and mallardite. It is used in industrial applications such as dyeing, porcelain glazing, and the manufacture of fertilizers and boiling oils.¹

In biochemistry, manganese is found in various superoxide dismutases.^{2,3} MnSO_4 is used as a source of manganese ion in biological research, such as in culturing of *Bacillus licheniformis* and the induction of chromosomal abnormalities in plants.^{4,5} MnSO_4 has been utilized to investigate the enzyme dependent glycosylation of endogenous glycoproteins in human skeletal muscle.⁶

A protocol that uses MnSO_4 in the removal of contaminating genomic DNA in RNA samples for RT-PCR has been published.⁷

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

This product is soluble in water (77 mg/ml), yielding a clear, light pink solution.

References

1. The Merck Index, 12th ed., Entry# 5782.
2. Martin, M. E., et al., A *Streptococcus mutans* superoxide dismutase that is active with either manganese or iron as a cofactor. *J. Biol. Chem.*, **261(20)**, 9361-9367 (1986).
3. Matsumoto, T., et al., Iron- and manganese-containing superoxide dismutases from *Methylobionas* J: identity of the protein moiety and amino acid sequence. *Biochemistry*, **30(13)**, 3210-3216 (1991).
4. Cromwick, A. M., and Gross, R. A., Effects of manganese (II) on *Bacillus licheniformis* ATCC 9945A physiology and γ -poly(glutamic acid) formation. *Int. J. Biol. Macromol.*, **17(5)**, 259-267 (1995).
5. Mukhopadhyay, M. J., and Sharma, A., Comparison of different plants in screening for Mn clastogenicity. *Mutat. Res.*, **242(2)**, 157-161 (1990).
6. Jiao, Y., et al., Manganese sulfate-dependent glycosylation of endogenous glycoproteins in human skeletal muscle is catalyzed by a nonglucose 6-P-dependent glycogen synthase and not glycogenin. *Biochim. Biophys. Acta*, **1427(1)**, 1-12 (1999).
7. Wang, G., et al., Bacterial DNA decontamination for reverse transcription polymerase chain reaction (RT-PCR). *J. Microbiol. Methods*, **51(1)**, 119-121 (2002).

GCY/RXR 5/03

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.