

UM-SCC-47 Squamous Carcinoma Cell Line

Cancer Cell Line
Cat. # SCC071

Pack size: $\geq 1 \times 10^6$
viable cells/vial

FOR RESEARCH USE ONLY
NOT FOR USE IN DIAGNOSTIC PROCEDURES
NOT FOR HUMAN OR ANIMAL CONSUMPTION

Store in liquid nitrogen



Certificate of Analysis

page 1 of 3

Background

Head and neck squamous-cell carcinoma (HNSCC) is the 6th most common type of cancer world-wide. The cancer may occur in the lip, oral cavity, nasal cavity, paranasal sinuses, salivary glands, pharynx and larynx. Risk factors include smoking, alcohol consumption, betel nut chewing, wood dust exposures and human papilloma virus (HPV) infections. Approximately 15% of HNSCC contain genomic DNA from HPV⁽²⁾. In particular, HPV-16 occurs in 90-95% of all HPV-positive HNSCC cases⁽²⁾. In oral and pharynx cancer, HPV DNA is found in the tonsils in 45-67% of the cases. HPV DNA is found in 13-25% of the cases in the hypopharynx, 12-18% in the oral cavity and 3-7% in the larynx⁽²⁾.

UM-SCC-47 is a unique head and neck squamous carcinoma cell line isolated from the primary tumor of the lateral tongue of a male⁽¹⁾ patient. The cell line contains 15-18 copies of integrated HPV-16 and approximately 10% of ALDH⁺ cancer stem cells.

STR Profile

| | |
|-----------------|------------------|
| D3S1358: 15 | D16S539: 8, 13 |
| TH01: 7, 9.3 | CSF1P0: 11, 13 |
| D21S11: 29, 30 | Penta D: 9, 10 |
| D18S51: 18 | vWA: 18 |
| Penta E: 12, 13 | D8S1179: 15 |
| D5S818: 11, 12 | TPOX: 10, 11 |
| D13S317: 8, 11 | FGA: 23, 25 |
| D7S820: 11 | Amelogenin: X, Y |

Cancer cell lines are inherently genetically unstable. Genetic instability may arise in the form of loss of heterozygosity of alleles at one or more genetic sites with increased passages.

Source

UM-SCC-47 was established at the University of Michigan⁽¹⁾ with written informed consent obtained from the patient and with the approval of the study by the Medical School Institutional Review Board as described by Brenner et al.

Quality Control Testing

- Each vial contains $\geq 1 \times 10^6$ viable cells.
- Cells are tested by PCR and are positive for HPV-16 and negative for Hepatitis A, B, C and HIV-1 & 2 viruses.
- Cells are negative for mycoplasma contamination.
- Each lot of cells are genotyped by STR analysis to verify the unique identity of the cell line.

Important Note



UM-SCC cell lines were derived in the lab of Dr. Thomas Carey at the University of Michigan and are exclusively distributed by Merck KGaA. PURCHASER may not distribute UM-SCC cells or derivatives to third parties.

THIS PRODUCT IS ONLY AVAILABLE FOR SALE TO ACADEMIC INSTITUTIONS OR NOT-FOR-PROFIT ENTITIES FOR USE UNDER THIS LIMITED USE LABEL LICENSE. FOR INFORMATION ON COMMERCIAL LICENSING OF UM-SCC-1 CELLS, INCLUDING LICENSING TO COMMERCIAL ENTITIES, PLEASE CONTACT licensing@emdmillipore.com.

Storage and Handling

UM-SCC-47 cells should be stored in liquid nitrogen. The cells can be cultured for at least 10 passages after the initial thaw without significantly affecting the cell marker expression and functionality.

Data



10% ALDH⁺ cancer stem cells detected in UM-SCC-47 cells by AldeRed 588-A (Cat. No. SCR150)

SPECIES LEGEND: H Human Ca Canine M Mouse R Rat Rb Rabbit B Bovine P Porcine WR Most Common Vertebrates

Please visit www.millipore.com for additional product information and references.

Submit your published journal article, and earn credit toward future purchases. Visit www.millipore.com/publicationrewards to learn more!

Protocols

Thawing of Cells

1. Do not thaw the cells until the recommended medium is on hand. Cells can grow on normal tissue culture ware surfaces without any additional coating.

Cells are thawed and expanded in DMEM High Glucose (EMD Millipore Cat. No. SLM-021-B), containing 10% FBS (EMD Millipore Cat. No. ES009-B) and Non-Essential Amino Acids (EMD Millipore Cat. No. TMS-001-C).

2. Remove the vial of frozen UM-SCC-47 cells from liquid nitrogen and incubate in a 37°C water bath. Closely monitor until the cells are completely thawed. Maximum cell viability is dependent on the rapid and complete thawing of frozen cells.

IMPORTANT: Do not vortex the cells.

3. As soon as the cells are completely thawed, disinfect the outside of the vial with 70% ethanol. Proceed immediately to the next step.
4. In a laminar flow hood, use a 1 or 2 mL pipette to transfer the cells to a sterile 15 mL conical tube. Be careful not to introduce any bubbles during the transfer process.
5. Using a 10 mL pipette, slowly add dropwise 9 mL of 10% FBS media (Step 1 above; pre-warmed to 37°C) to the 15 mL conical tube.

IMPORTANT: Do not add the entire volume of media all at once to the cells. This may result in decreased cell viability due to osmotic shock.

6. Gently mix the cell suspension by slowly pipetting up and down twice. Be careful not to introduce any bubbles.

IMPORTANT: Do not vortex the cells.

7. Centrifuge the tube at 300 x g for 2-3 minutes to pellet the cells.
8. Decant as much of the supernatant as possible. Steps 5-8 are necessary to remove residual cryopreservative (DMSO).
9. Resuspend the cells in a total volume of 10 % FBS medium (pre-warmed to 37°C).
10. Transfer the cell mixture into a T75 tissue culture flask.
11. Incubate the cells at 37°C in a humidified incubator with 5% CO₂.
12. The next day, exchange the medium with fresh 10% FBS media pre-warmed to 37°C. Exchange with fresh medium every two to three days thereafter.
13. When the cells are approximately 90% confluent, they can be dissociated with Accutase (EMD Millipore Cat. No. SCR005) or trypsin-EDTA (EMD Millipore Cat. No. SM-2003-C) and further passaged or, alternatively, frozen for later use.

Subculturing of Cells

1. Carefully remove the medium from the T75 tissue culture flask containing the confluent layer of UM-SCC-47 cells.
2. Apply 3-5 mL of Accutase or trypsin-EDTA solution and incubate in a 37°C incubator for 3-5 minutes.
3. Inspect the flask and ensure the complete detachment of cells by gently tapping the side of the flask with the palm of your hand.
4. Add 8 mL of 10% FBS medium (pre-warmed to 37°C) to the plate.
5. Gently rotate the flask to mix the cell suspension. Transfer the dissociated cells to a 15 mL conical tube.
6. Centrifuge the tube at 300 x g for 3-5 minutes to pellet the cells.
7. Discard the supernatant, then loosen the cell pellet by tapping the tip of the tube with a finger.
8. Apply 2 mL of 10% FBS media (pre-warmed to 37°C) to the conical tube and resuspend the cells thoroughly.

IMPORTANT: Do not vortex the cells.

9. Count the number of cells using a hemocytometer.
10. Plate the cells to the desired density (typical split ratio is 1:3 to 1:6).

Cryopreservation of Cells

UM-SCC-47 cells can be frozen in the expansion media plus 10% DMSO using a Nalgene slow freeze Mr. Frosty container.

References

1. Brenner, J.C., Graham, M.P., Kumar, B., Saunders, L.M., Kupfer, R., Lyons, R.H., Bradford, C.R., Carey, T.E. (2010) Genotyping of 73 UM-SCC head and neck squamous cell carcinoma cell lines. *Head Neck* 34 (4): 417-26.
2. Perez-Ordoñez, B., Beauchemin, M., and Jordan, R.C. (2006) Molecular biology of squamous cell carcinoma of the head and neck. *J. Clin Pathol* 59(5): 445-453.

■ antibodies ■ Multiplex products ■ biotools ■ cell culture ■ enzymes ■ kits ■ proteins/peptides ■ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

EMD Millipore Corporation, 28820 Single Oak Drive, Temecula, CA 92590, USA 1-800-437-7500

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502

FOR RESEARCH USE ONLY. Not for use in diagnostic procedures. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited.

Millipore®, the M mark, Upstate®, Chemicon®, Linco® and all other trademarks, unless specifically identified above in the text as belonging to a third party, are owned by Merck KGaA, Darmstadt. Copyright ©2008-2015 Merck KGaA, Darmstadt. All rights reserved.



We Buy 100% Certified Renewable Energy

LIMITED USE LABEL LICENSE

BY OPENING THIS PRODUCT, YOU (“PURCHASER”) HEREBY REPRESENT THAT YOU HAVE THE RIGHT AND AUTHORITY TO LEGALLY BIND YOURSELF AND/OR YOUR INSTITUTION, AS APPLICABLE, AND CONSENT TO BE LEGALLY BOUND BY ALL OF THE TERMS OF THIS LIMITED USE LABEL LICENSE. IF YOU DO NOT AGREE TO COMPLY WITH THESE TERMS, PLEASE DO NOT OPEN THE PRODUCT AND CALL EMD MILLIPORE (“SELLER”) CUSTOMER SERVICE (1-800-645-5476) TO ARRANGE TO RETURN THE PRODUCT FOR A REFUND.

SELLER hereby conveys to PURCHASER the non-exclusive and non-transferable right to use the purchased amount of the Product for non-commercial Research Purposes only.

“Research Purposes” means any biological research and development uses and specifically excludes the following uses of whatever kind or nature:

- use of UM-SCC-47 or derivatives in the development, testing, or manufacture of a Commercial Product
- use of UM-SCC-47 Cells or derivatives in a Commercial Product
- sale or licensing of UM-SCC-47 Cells or derivatives
- clinical or therapeutic applications including the production of materials for clinical trials
- use in or administration to humans

“Commercial Product” means any product intended for sale, or for drug, diagnostic, or therapeutic use.

Access to UM-SCC-47 Cells is limited solely to those officers, employees and students of PURCHASER’s institution who need access to perform research. PURCHASER may not distribute UM-SCC-47 Cells or derivatives to third parties. PURCHASER shall comply with all applicable laws in its use and handling of the Product and shall keep it under reasonable safe and secure conditions to prevent unauthorized use or access.

This AGREEMENT will remain in effect for as long as PURCHASER possesses the Product.

THIS PRODUCT IS ONLY AVAILABLE FOR SALE TO ACADEMIC INSTITUTIONS OR NOT-FOR-PROFIT ENTITIES FOR USE UNDER THIS LIMITED USE LABEL LICENSE. FOR INFORMATION ON COMMERCIAL LICENSING OF UM-SCC-47 CELLS, INCLUDING LICENSING TO COMMERCIAL ENTITIES, PLEASE CONTACT licensing@emdmillipore.com.

■ antibodies ■ Multiplex products ■ biotools ■ cell culture ■ enzymes ■ kits ■ proteins/peptides ■ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

EMD Millipore Corporation, 28820 Single Oak Drive, Temecula, CA 92590, USA 1-800-437-7500

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502

FOR RESEARCH USE ONLY. Not for use in diagnostic procedures. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited.

Millipore®, the M mark, Upstate®, Chemicon®, Linco® and all other trademarks, unless specifically identified above in the text as

belonging to a third party, are owned by Merck KGaA, Darmstadt. Copyright ©2008-2015 Merck KGaA, Darmstadt. All rights reserved.



We Buy 100% Certified
Renewable Energy