

## Product Information

**PDGFR  $\alpha$  (T674I), active, GST-tagged, human PRECISIO® Kinase recombinant, expressed in *Sf9* cells**

Catalog Number **SRP5304**  
Storage Temperature  $-70\text{ }^{\circ}\text{C}$

Synonyms: CD140A, PDGFR2, MGC74795, Rhe-PDGFR A

### Product Description

PDGFR  $\alpha$  (platelet-derived growth factor receptor  $\alpha$ ) is a member of the PDGFR family of membrane receptors with intrinsic tyrosine kinase activity. Aberrant expression of PDGFR  $\alpha$  has been linked to developmental abnormalities in vertebrate models and has been implicated in multiple disease states in humans. There is widespread expression of PDGFR  $\alpha$  in renal cell types involved in fibrotic and sclerosing processes.<sup>1</sup> PDGF and its receptor PDGFR  $\alpha$  are inducers of fibrosis in the repair phase of inflammatory bowel disease and they may also be involved in the active inflammatory phase.<sup>2</sup> PDGFR  $\alpha$  (T674I) is one of the mutant forms of PDGFR  $\alpha$ .

Recombinant human PDGFR  $\alpha$  (T674I) (550-end) was expressed by baculovirus in *Sf9* insect cells using an N-terminal GST-tag. The gene accession number is NM\_006206. It is supplied in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, and 25% glycerol.

Molecular mass:  $\sim$ 95 kDa


### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.


### Storage/Stability

The product ships on dry ice and storage at  $-70\text{ }^{\circ}\text{C}$  is recommended. After opening, aliquot into smaller quantities and store at  $-70\text{ }^{\circ}\text{C}$ . Avoid repeated handling and multiple freeze/thaw cycles.

**Figure 1.**  
SDS-PAGE Gel of Typical Lot:  
 $\geq 70\%$  (SDS-PAGE, densitometry)



**Figure 2.**  
Specific Activity of Typical Lot:  
16–24 nmole/min/mg



### Procedure

#### Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 20 mM  $\text{MgCl}_2$ , 25 mM  $\text{MnCl}_2$ , 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/ $\mu\text{L}$  BSA solution.

Kinase Solution – Dilute the active PDGFR  $\alpha$  (T674I) (0.1  $\mu\text{g}/\mu\text{L}$ ) with Kinase Dilution Buffer to the desired concentration.

Note: The specific activity plot may be used as a guideline (see Figure 2). It is recommended the researcher perform a serial dilution of active PDGFR  $\alpha$  (T674I) kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 mL of Kinase Assay Buffer. Store in 200  $\mu\text{L}$  aliquots at  $-20\text{ }^{\circ}\text{C}$ .

$\gamma$ - $^{33}\text{P}$ -ATP Assay Cocktail (250  $\mu\text{M}$ ) – Combine 5.75 mL of Kinase Assay Buffer, 150  $\mu\text{L}$  of 10 mM ATP Stock Solution, 100  $\mu\text{L}$  of  $\gamma$ - $^{33}\text{P}$ -ATP (1 mCi/100  $\mu\text{L}$ ). Store in 1 mL aliquots at  $-20\text{ }^{\circ}\text{C}$ .

Substrate Solution – MBP Protein substrate diluted in distilled water to a final concentration of 1 mg/mL.

1% phosphoric acid solution – Dilute 10 mL of concentrated phosphoric acid to a final volume of 1 L with water.

#### Kinase Assay

This assay involves the use of the  $^{33}\text{P}$  radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the active PDGFR  $\alpha$  (T674I), Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The  $\gamma$ - $^{33}\text{P}$ -ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20  $\mu\text{L}$ :
  - 10  $\mu\text{L}$  of Kinase Solution
  - 5  $\mu\text{L}$  of Substrate Solution
  - 5  $\mu\text{L}$  of cold water ( $4\text{ }^{\circ}\text{C}$ )
3. Set up a blank control as outlined in step 2, substituting 5  $\mu\text{L}$  of cold water ( $4\text{ }^{\circ}\text{C}$ ) for the Substrate Solution.
4. Initiate each reaction with the addition of 5  $\mu\text{L}$  of the  $\gamma$ - $^{33}\text{P}$ -ATP Assay Cocktail, bringing the final reaction volume to 25  $\mu\text{L}$ . Incubate the mixture in a water bath at  $30\text{ }^{\circ}\text{C}$  for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20  $\mu\text{L}$  of the reaction mixture onto an individually precut strip of phosphocellulose P81 paper.

6. Air dry the precut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of  $\sim 10$  minutes each.
7. Set up a radioactive control to measure the total  $\gamma$ - $^{33}\text{P}$ -ATP counts introduced into the reaction. Spot 5  $\mu\text{L}$  of the  $\gamma$ - $^{33}\text{P}$ -ATP Assay Cocktail on a precut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

#### Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$\text{SR} = \frac{\text{cpm of } 5\ \mu\text{L of } \gamma\text{-}^{33}\text{P-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)  
nmole – 1.25 nmole (5  $\mu\text{L}$  of 250  $\mu\text{M}$  ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta\text{cpm} \times (25/20)}{\text{SR} \times E \times T}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

$\Delta\text{cpm}$  = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

#### References

1. Floege, J. et al., Expression of PDGF alpha-receptor in renal arteriosclerosis and rejecting renal transplants. *J. Am. Soc. Nephrol.*, **9**(2), 211-23 (1998).
2. Kumagai, S. et al., Platelet-derived growth factor and its receptors are expressed in areas of both active inflammation and active fibrosis in inflammatory bowel disease. *Tohoku J. Exp. Med.*, **195**(1), 21-33 (2001).

PRECISIO is a registered trademark of Sigma-Aldrich Co. LLC.

RC,MAM 12/12-1