

Product Information

Anti-Profilin 1 (C-terminal)

produced in rabbit, affinity isolated antibody

Catalog Number **P7624**

Product Description

Anti-Profilin 1 (C-terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to amino acid residues 128-140 of human profilin 1, conjugated to KLH. The corresponding sequence is identical in rat and mouse. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-Profilin 1 (C-terminal) recognizes human, mouse, and rat profilin 1. Applications include immunoblotting (~15 kDa), immunoprecipitation, and immunofluorescence. Detection of the profilin 1 band by immunoblotting is specifically inhibited by the immunizing peptide.

Profilin 1 is a ubiquitous actin monomer-binding protein involved in actin polymerization in response to extracellular signals.¹ Three profilin genes have been identified: profilin 1, 2, and 3. Profilin 1 is the most ubiquitous and abundant, and is highly expressed throughout development and adulthood in most tissues including brain. Profilin 2 is the neuronal specific isoform and profilin 3 is a testis specific isoform.^{2,3} Profilins of eukaryotic cells are small (12-15 kDa) cytoplasmic proteins that bind to actin monomers, polyphosphoinositides and polymers of L-proline.¹ Profilins were shown to be important for normal cell proliferation, differentiation and motility.⁴ Deletion of profilin 1 gene leads to an embryonic lethal phenotype.⁵

Profilin 1 is a potent regulator of actin filament dynamics. Although profilin 1 prevents spontaneous actin polymerization by complexing with unpolymerized actin *in vivo*, actin-profilin complexes can be added to free barbed ends, thereby stimulating actin polymerization. When bound to actin, profilin functions as an ATP nucleotide exchange factor recharging ADP-actin with ATP.^{6,7} Dissociation of the profilin-actin complex is caused by binding of profilin to phosphatidylinositol 4,5-bisphosphate (PIP₂), which liberates actin for polymerization.⁸ Profilin binds with high affinity to poly-L-proline stretches found in many cellular proteins associated with the cytoskeleton.

Profilin is recruited to sites of active cytoskeletal assembly through its interaction with proteins such as N-WASP, the ARP2/3 complex, p140mDia, VASP and Mena.⁷ The interactions of profilin 1 with actin, proline-rich proteins and PIP₂ influences neuronal differentiation of PC12 cells.⁹ Profilin 1 was suggested to act as a tumor suppressor protein based on its reduced expression in several types of invasive cancers and its ability to suppress tumorigenicity when over-expressed in breast cancer cells.^{4,10} Deletion of profilin 1 is associated with Miller-Dieker syndrome.¹¹

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1 mg/mL

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working concentration of 1-2 µg/mL is recommended using a whole extract of human HeLa cells, applying a chemiluminescent detection reagent.

Immunoprecipitation: 5-10 µg of the antibody immunoprecipitates profilin 1 from mouse NIH-3T3 cell lysates.

Indirect immunofluorescence: a working concentration of 10-20 µg/mL is recommended using rat NRK cells.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Goldschmidt-Clermont, P.J., et al., *J. Cell Biol.*, **113**, 1081-1089 (1991).
2. Honore, B., et al., *FEBS Lett.*, **330**, 151-155 (1993).
3. Braun, A., et al., *Gene*, **283**, 219-225 (2002).
4. Wittenmayer, N., et al., *Mol. Biol. Cell*, **15**, 1600-1608 (2004).
5. Witke, W., et al., *Proc. Natl. Acad. Sci. USA*, **98**, 3832-3836 (2001).
6. Goldschmidt-Clermont, P.J., et al., *Mol. Biol. Cell*, **3**, 1015-1024 (1992).
7. Suetsugu, S., et al., *EMBO J.*, **17**, 6516-6526 (1998).
8. Lassing, I., and Lindberg, U., *Nature*, **314**, 472-474 (1985).
9. Lambrechts, A., et al., *J. Cell Sci.*, **119**, 1570-1578 (2006).
10. Roy, P., and Jacobson, K., *Cell Motil. Cytoskel.*, **57**, 84-95 (2004).
11. Kwiatkowski, D.J., et al., *Am. J. Hum. Genet.*, **46**, 559-567 (1990).

ST,KAA,PHC 10/06-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.