

Product Information

Anti-Lamin A/C R453W antibody, Mouse monoclonal

Clone 12A-2F5, purified from hybridoma cell culture

SAB4200421

Product Description

Anti-Lamin A/C R453W (mouse IgG1 isotype) is derived from the hybridoma 12A-2F5 produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with a synthetic peptide containing the R453W substitution of human Lamin A/C (GeneID: 4000).¹ The isotype is determined by ELISA using Mouse Monoclonal Antibody Isotyping Reagents (Cat. No. ISO2). The antibody is purified from culture supernatant of hybridoma cells grown in a bioreactor.

Anti-Lamin A/C R453W recognizes a human Lamin A/C recombinant protein containing the R453W substitution. This antibody does not recognize the Lamin A/C wild-type protein. The antibody may be used in several immunochemical techniques including immunoblotting, immunoprecipitation and immunofluorescence.¹

Lamin A is a structural protein of the nuclear lamina. The nuclear lamina is a meshwork of intermediate filaments that underlies the inner face of the nuclear envelope.² The major components of the nuclear lamina are the lamins that may be classified into two types, A and B. Both A- and B- type lamins are characterized by an α -helical rod domain to enable assembly into filaments, a nuclear localization sequence, and a carboxy-terminal CAAX box isoprenylation sequence for nuclear membrane targeting.³ A-type lamins, lamin A and lamin C, are products of a single gene, LMNA, which are produced by alternative splicing, resulting in proteins of 664 and 572 amino acids, respectively. ⁴ The first 566 amino acids of Lamin A and C are identical. Prelamin A, the precursor of Lamin A, has 98 unique amino acids and is farnesylated at its carboxy terminus after synthesis. The last 18 amino acids, which contain the farnesyl group, are removed by an endoproteolytic cleavage, producing the mature Lamin A.⁵

Lamins are expressed in most somatic cells. They interact with integral proteins of the inner membrane of the nuclear envelope, such as LAPs 1 and 2 (lamina-associated polypeptides), LBR (Lamin B receptor) and emerin.⁶ They also interact with chromatin and nuclear pore complexes.⁵

Lamin A is cleaved into a 47 kDa fragment during apoptosis. This cleavage seems to be essential for chromatin condensation and nuclear disassembly in apoptosis.^{3,7} Mutations in Lamin A and C have been linked to a variety of rare human diseases including muscular dystrophy, lipodystrophy, cardiomyopathy, neuropathy and progeroid syndromes (collectively termed laminopathies) and to premature aging (Hutchinson-Gilford progeria syndrome).^{8,9} Most diseases arise from dominant, missense mutations. Lamin A/C R453W antibody is a critical tool for basic and clinical research of the mutant A-type lamins in primary patient cells.

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody Concentration: ~ 1.0 mg/mL

Precautions and Disclaimer

This product is for research use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze at -20 °C in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 0.2-0.5 µg/mL is recommended using human Lamin A/C partial recombinant protein containing the R453W substitution.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Roblek, M., et al., *PLoS ONE*, **5**, e10604 (2010).
2. Aebi, U., et al., *Nature*, **323**, 560-564 (1986).
3. Rao, L., et al., *J. Cell Biol.*, **135**, 1441-1455 (1996).
4. Lloyd, D.J., et al., *Hum. Mol. Gen.*, **11**, 769-777 (2002).
5. Worman, H.J., and Courvalin, J.C., *Trends Cell Biol.*, **12**, 591-598 (2002).
6. Schirmer, E.C., et al., *J. Cell Biol.*, **153**, 479-489 (2001).
7. Ruchaud, S., et al., *EMBO J.*, **21**, 1967-1977 (2002).
8. Worman, H.J., and Courvalin, J.C., *J. Clin. Invest.*, **113**, 349-351 (2004).
9. Plasilova, M., et al., *J. Med. Genet.*, **41**, 609-614 (2004).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

SAB4200421dat Rev 06/21

For research use only. Not for use in diagnostic procedures.