



3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

18S + 28S Ribosomal RNA From Calf Liver

Product No. **R 0889**

Lot 063K0353
Store at -70 °C

PRODUCT SUMMARY

Suitable for use as a molecular weight marker for formaldehyde agarose gel electrophoresis.

Concentration: 2.0 mg/ml

STORAGE BUFFER

10 mM Tris, pH 8.0
1 mM EDTA

10X MOPS ELECTROPHORESIS BUFFER

400 mM MOPS
100 mM Sodium Acetate
10 mM EDTA

LOADING DYE

50 % (v/v) glycerol
1 mM EDTA
0.4 % (w/v) Bromophenol Blue
0.4 % (w/v) Xylene Cyanol

SUITABILITY ASSAY

E. coli 18S + 28S ribosomal RNA (rRNA) sample solutions were prepared for electrophoresis as follows:

1.0 µg 18S + 28S Ribosomal RNA
2.5 µl 10X Running Buffer
3.5 µl 37% Formaldehyde
10 µl Deionized Formamide
1 µl Ethidium Bromide (0.2 mg/ml)

SUITABILITY ASSAY(continued)

The above sample solution was incubated at 65 °C for 10 minutes and immediately cooled on ice. 2 µl of loading dye was added and the entire sample solution (1 µg 18S + 28S Ribosomal RNA) was run along with appropriate RNA markers on a denaturing (formaldehyde) agarose gel. Electrophoresis was performed in a submarine-type apparatus at 100 volts for 2 hours in 1X MOPS electrophoresis buffer with buffer recirculation. Two bands were resolved, and the sizes were consistent with 18S and 28S ribosomal RNA.

Fragment Sizes:

18S rRNA approx. 2000 bases
28S rRNA approx. 5300 bases

References:

1. Sambrook, J., et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory (1989), p.202.
2. Fasman, G.D., ed., Practical Handbook of Biochemistry and Molecular Biology, CRC Press, (1986), p.464.

6/03