

Product Information

PHOSPHOSERINE ANTIBODY INHIBITOR

Product Number **P 3717**

Product Description

Phosphoserine Antibody Inhibitor blocks specifically the reactivity of anti-phosphoserine antibody, but not that of anti-phosphothreonine- or anti-phosphotyrosine-specific antibodies. The product is useful in immunoblotting and in ELISA.

Phosphoserine Antibody Inhibitor may be used for the specific inhibition of the reactivity of anti-phosphoserine antibodies using various immunochemical assays such as immunoblotting and ELISA

Antibodies reacting specifically with phosphorylated amino acid residues are useful as analytical and preparative tools and serve in the identification, quantification and immunoaffinity isolation of activated cellular proteins. Often it is desirable to verify the specificity of the antibodies for the particular phosphorylated amino acid residue. Unfortunately, the use of the appropriate phosphorylated amino acid to specifically block the reactivity of the antibodies, does not fulfill the requirement, due to the small size of the blocking amino acid. The availability of a preparation of O-Phospho-L-Serine conjugated to BSA carrier should aid in this task.

Reagents

Phosphoserine Antibody Inhibitor is a solution of O-Phospho-L-Serine conjugated to BSA, in 0.01 M phosphate buffered saline pH 7.4, with 15 mM sodium azide as a preservative.

Precautions and Disclaimer

Due to the sodium azide content a material safety sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for a maximum of one month. For extended storage freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Product Profile

A working dilution of 2 µg/ml is determined by the specific inhibition of the reactivity of Monoclonal Anti-Phosphoserine (Product No. P 3430), used in immunoblotting of rat cerebral cortex extract, and in ELISA.

Notes

1. In order to obtain best results in different techniques and preparations we recommend each user determine optimal working dilutions by titration test.
2. Decreasing or increasing the dilution of the reactive antibody will affect the amount of the inhibitor required to achieve the desired degree of blocking .

JWM/DAA 11/2002

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.