

## Product Information

**Anti-TNFAIP1**  
antibody produced in chicken,  
affinity isolated antibody

Catalog Number **GW22047**  
Formerly listed as GenWay Catalog Number 15-288-22047,  
Tumor necrosis factor, alpha-induced protein 1, endothelial  
Antibody.

Storage Temperature Store at -20 °C

Synonyms: Tumor necrosis factor, alpha-induced protein 1,  
B12 protein

### Product Description

This protein was identified as a gene whose expression can be induced by the tumor necrosis factor alpha (TNF) in umbilical vein endothelial cells. Studies of a similar gene in mouse suggest that the expression of this gene is developmentally regulated in a tissue-specific manner.

NCBI Accession number: NP\_066960.1  
Swiss Prot Accession number: Q13829

Gene Information: Human .. TNFAIP1 (7126)

Immunogen: Recombinant protein Tumor necrosis factor,  
alpha-induced protein 1

Immunogen Sequence: GI # 10863937, sequence 1 - 316

The product is a clear, colorless solution in phosphate buffered saline, pH 7.2, containing 0.02% sodium azide.

Species Reactivity: Human, mouse, rat

Tested Applications: ELISA, WB

Recommended Dilutions: Recommended starting dilution for Western blot analysis is 1:500, for tissue or cell staining 1:200.

Note: Optimal concentrations and conditions for each application should be determined by the user.

### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Due to the sodium azide content a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

### Storage/Stability

For continuous use, store at 2–8 °C for up to one week. For extended storage, store in -20 °C freezer in working aliquots. Repeated freezing and thawing, or storage in "frostfree" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

LPG,MAM12/09-1