

53876 Atto 647N alkyne

Application

Atto 647N belongs to a new generation of fluorescent labels for the red spectral region. The dye is designed for application in the area of life science, e.g. labeling of DNA, RNA or proteins. Characteristic features of the label are strong absorption, excellent fluorescence quantum yield, high photostability, excellent ozone resistance, good solubility, and very little triplet formation. Atto 647N is a cationic dye. After coupling to a substrate the dye carries a net electrical charge of $^{+1}$.


In common with most Atto-labels, absorption and fluorescence are independent of pH in the range of 2 to 11, used in typical applications. As supplied Atto 647N consists of a mixture of two isomers with practically identical absorption and fluorescence properties.

The **alkyne** modification is used in the Huisgen reaction ("Click Chemistry").

Product Description

| | |
|---------------------|--|
| MW | 783 g/mol |
| λ_{abs} | 646 nm |
| ε_{max} | $1.5 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ |
| λ_{fl} | 664 nm |
| η_{fl} | 65 % |
| τ_{fl} | 3.5 ns |
| CF_{260} | 0.04 |
| CF_{280} | 0.03 |

Optical data of the carboxy derivative (in aqueous solution)



Storage: Store at -20°C and protected from light.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

The vibrant M and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. Detailed information on trademarks is available via publicly accessible resources. © 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

